proLékaře.cz | 13.11.2026

2. Havrdová E. Roztroušená skleróza. Cesk Slov Neurol N 2008; 71/104(2): 121–132.
3. Aronson KJ. Quality of life among persons with multiple sclerosis and their caregivers. Neurology 1997; 48(1): 74–80.
4. Kalb R. Multiple Sclerosis: a model of Psychosocial Support. London: National Multiple Sclerosis Society 2000.
5. Motl RW, McAuley E, Snook EM, Gliottoni RC. Physical activity and quality of life in multiple sclerosis: intermediary roles of disability, fatigue, mood, pain, self-efficacy and social support. Psychol Health Med 2009; 14(1): 111–124. doi: 10.1080/13548500802241902.
6. José Sá M. Psychological aspects of multiple sclerosis. Clin Neurol Neurosurg 2008; 110(9): 868–877.
7. Gibson J, Frank A. Supporting individuals with disabling multiple sclerosis. J R Soc Med 2002; 95(12): 580–586.
8. Mohr DC, Likosky W, Bertagnolli A, Goodkin DE, Van Der Wende J, Dwyor P et al. Telephone-administered cognitive-behavioral therapy for the treatment of depressive symptoms in multiple sclerosis. J Consult Clin Psychol 2000; 68(2): 356–361.
9. Tesar N, Baumhackl U, Kopp M, Günther V. Effects of psychological group therapy in patients with multiple sclerosis. Acta Neurol Scand 2003; 107(6): 394–399.
10. Langenmayr A, Schöttes N. Psychotherapy with multiple-sclerosis patients. Psychol Rep 2000; 86(2): 495–508.

