

# The Changing Microbiological Pattern in Patients with Confirmed Bacterial Meningitis after Post-craniotomy Surgery

Změna mikrobiologického vzorku u pacientů s potvrzenou bakteriální meningitidou po postkraniotomiální operaci

## Abstract

Post-craniotomy meningitis (PCM) is an important challenge in neurosurgery. In this study, PCM was diagnosed in 22 of the 4,392 post-craniotomy patients between January 1, 2008 and December 31, 2012. There were significant statistical differences between the seven patients with *Acinetobacter baumannii* meningitis and 15 patients with non-*Acinetobacter baumannii* meningitis with respect to the history traumatic head injury (OR = 5.873, 95% CI 1.575–17.511;  $p = 0.004$ ) and infected sites other than the central nervous system (OR = 0.872, 95% CI 0.665–1.621;  $p = 0.003$ ). We provide the most up-to-date insight to physicians on the changing pattern of infectious agents in PCM.

## Souhrn

Meningitida po kraniotomii (PCM) představuje důležitou výzvu v neurochirurgii. Tato práce popisuje diagnostikovanou PCM u 22 z 4 392 pacientů s provedenou kraniotomií v intervalu mezi 1. lednem 2008 a 31. prosincem 2012. U sedmi pacientů s meningitidou způsobenou bakterií *Acinetobacter baumannii* a u 15 pacientů s neprokázanou *Acinetobacter baumannii* meningitidou byly signifikantní rozdíly ve výskytu traumatu hlavy (OR = 5,873, 95% CI 1,575–17,511;  $p = 0,004$ ), a lokalizací infekce mimo centrální nervový systém (OR = 0,872, 95% CI 0,665–1,621;  $p = 0,003$ ). Naše sdělení nabízí lékařům aktuální pohled na měnící se vzorec infekčních agensů u PCM.

The present work was partially supported by a grant from the Changhua Christian Hospital (grant 103-CCH-IPR-001 and 104-CCH-IPR-001). The authors thank the staff of the Clinical Microbiology Laboratory, Department of Medical Records, and Department of Computing of Changhua Christian Hospital for case findings. The authors thank prof. Min-Chi Liu for proof reading of this article. This research project would not have been possible without the support of many people. The authors wish to express their gratitude to the staff of the Infection Control Committee, Division of Infectious Diseases, Department of Nursing, Department of Healthcare Quality, and Department of Pharmacology of the Changhua Christian Hospital who were extremely helpful and provided invaluable assistance and support. We thank Fu-Chou Chen serving at the Taichung Veterans General Hospital for reviewing this article.

The authors declare they have no potential conflicts of interest concerning drugs, products, or services used in the study.

Autoři deklarují, že v souvislosti s předmětem studie nemají žádné komerční zájmy.

The Editorial Board declares that the manuscript met the ICMJE "uniform requirements" for biomedical papers.

Redakční rada potvrzuje, že rukopis práce splnil ICMJE kritéria pro publikace zasílané do biomedicínských časopisů.

Ch. Chang-Hua<sup>1</sup>, Y. Hua-Cheng<sup>2</sup>, Ch. Yu-Jun<sup>3</sup>, L. Chih-Wei<sup>4</sup>

Changhua Christian Hospital, Changhua, Taiwan:

<sup>1</sup> Division of Infectious Disease, Department of Internal Medicine

<sup>2</sup> Division of Neurosurgery, Department of Surgery

<sup>3</sup> Laboratory of Epidemiology and Biostatistics

<sup>4</sup> Department of Medical Imaging



Chen Chang-Hua, MD

Division of Infectious Disease  
Department of Internal Medicine  
Changhua Christian Hospital  
135 Nanshao Street  
Changhua 500  
Taiwan  
e-mail: 76590@cch.org.tw

Accepted for review: 24. 6. 2016

Accepted for print: 18. 8. 2016

## Key words

meningitis – neurosurgery – *Acinetobacter baumannii* – *Staphylococcus aureus* – head trauma

## Klíčová slova

meningitida – neurochirurgie – *Acinetobacter baumannii* – *Staphylococcus aureus* – poranění hlavy

## Introduction


Post-craniotomy meningitis (PCM) results in severe morbidity and mortality [1,2]. The rates of PCM have been reported to vary from 0.3 to 10%, and *Staphylococcus aureus* (*S. aureus*) is the major pathogen [2,3]. Some studies revealed that the rate of PCM due to *Acinetobacter baumannii* complex (*A. baumannii*) is increasing [1,4–6], and *A. baumannii* meningitis (ABM) causes high mortality

rates [5]. In addition, the rate of PCM due to *A. baumannii* has become accretive from China, Iran and Turkey [1,4–6]. The limited epidemiological data can be perplexing for physicians, in both making accurate diagnosis and choosing effective antibiotics, hence an urgent clinical need was behind the motivation to perform this study to determine the current pattern of infectious agents of PCM in Changhua Christian Hospital (CCH).

## Observation

The study was approved by the ethics committee of CCH (CCH IRB No. 131118). CCH is a 1,800-bed tertiary referral medical center, and more than 1,500 craniotomies are performed in CCH every year. This retrospective study included 22 patients who developed post-neurosurgical meningitis among the 4,392 patients who underwent craniotomy procedures between January 1, 2008 and December 31, 2012. Cases of post-craniotomy meningitis were identified from microbiological databases and medical records according to the International Classification of Diseases, ninth edition, clinical modifications (ICD-9-CM) at CCH in central Taiwan. The diagnosis of meningitis was based on the inclusion criteria and included both significant clinical evidence and microbiological confirmation. The definition of surgical site infection (SSI) is according to CDC/NHSN surveillance definitions [7]. The Vitek-2 System (BioMérieux, Marcy l'Etoile, France) was used to identify the phenotype of isolates. SPSS version 21.0 for Windows (SPSS Inc., Chicago, IL, USA) were used statistical analysis.

During the study period, PCM was diagnosed in 22 of the 4,392 operated patients. The prevalence of PCM in CCHS was  $134.4 \pm 76.8$  per 100,000 patient-years during the study period. The incidence of PCM was 0.5% among the 4,392 operated patients in CCH. The five-year longitudinal analysis of healthcare-associated infections (HAI), SSI, and total number of craniotomies and total number of PCMs during the study period is shown in Graph 1. The key risk factors among the 4,392 patients who underwent post-craniotomy operation and



**Graph 1.** The correlation among healthcare associated infection, surgical site infection, and total craniotomy number and total post-craniotomy meningitis during the study period was shown.

Others included *P. aeruginosa* in 2 patients, *K. pneumoniae* in 2 patients, *E. cloacae* in one patient, *E. coli* in 1 patient, and *E. faecalis* in 1 patient.

HAI – healthcare associated infection, SSI – surgical site infection.

**Tab. 1.** Demography and clinical characteristics between *A. baumannii* and non-*A. baumannii* patients who happened post-craniotomy meningitis.

Parameters		Total (n = 22)	Meningitis		p <sup>1</sup>	OR (95% CI)	p <sup>2</sup>
			Ab (n = 7)	non-Ab (n = 15)			
infected sites other than the CNS	yes	14 (63.6)	1 (14.3)	13 (86.7)	0.002	0.665–1.621	0.003
	no	8 (36.4)	6 (84.7)	2 (13.3)			
head trauma	yes	6 (27.3)	5 (71.4)	1 (6.7)	0.004	5.873 (1.575–17.511)	0.004
	no	16 (72.7)	2 (28.6)	14 (93.3)			

<sup>1</sup> p value between those with meningitis and those without meningitis.

<sup>2</sup> p value for odds ratio (OR).

Data presents with number with percentage (parenthesis), n (%). The non-Ab group included *S. aureus* in 8 patients, *P. aeruginosa* in 2 patients, *K. pneumoniae* in 2 patients, *E. cloacae* in 1 patient, *E. coli* in one patient, and *E. faecalis* in 1 patient.

Ab – *Acinetobacter baumannii* complex, CNS – central nerve system.

Tab. 2. Demography and clinical characteristics of seven patients who happened post-craniotomy *A. baumannii* meningitis.

Patient characteristics	Patient 1	Patient 2	Patient 3	Patient 4	Patient 5	Patient 6	Patient 7
age (years)/sex/FATD (days)	41/M/8	32/M/30	65/M/8	56/M/10	41/M/5	38/F/12	75/F/2
<b>Neurosurgical diagnosis</b>							
head traumatic injury/intracerebral hemorrhage/ /intracerebral tumor	Y/N/N	Y/N/N	N/Y/N	Y/N/N	Y/N/N	N/N/Y	Y/N/N
<b>co-morbidity</b>							
diabetes mellitus/malignancy/coronary artery disease	N/N/N	N/N/N	Y/N/N	Y/N/N	N/N/N	N/N/N	Y/N/N
other source of <i>Acinetobacter</i>	blood	no	tracheal	tracheal	tracheal	N	N
<b>Risk Factors</b>							
emergency procedure/CSF leak/preoperative use of corticoids/prophylactic antibiotics before operation	Y/Y/N/N	N/N/N/Y	Y/Y/N/N	Y/Y/N/N	N/N/Y/N	Y/Y/N/N	Y/Y/N/Y
tracheal intubation/tracheotomy/mechanical ventilation	Y/N/Y	Y/N/Y	Y/N/Y	Y/N/Y	N/N/Y	Y/N/Y	Y/Y/Y
external ventricular drain/central venous catheters inser- tion/drain placement > 72 hours	Y/Y/N	N/N/N	Y/N/N	Y/Y/N	Y/N/N	Y/N/Y	Y/Y/Y
surgery duration (h) > 4.5/length of preop stay > 2 days/ /ICU admission	N/N/Y	Y/N/N	N/N/Y	N/N/Y	N/N/N	N/N/Y	Y/Y/Y
urinary catheterization/on NG tube/repeat operations	Y/Y/N	N/Y/N	Y/Y/N	Y/Y/N	N/N/N	Y/Y/N	Y/Y/Y
head trauma/hydrocephalus/use of perioperative antibiotics	Y/N/N	Y/N/Y	N/N/N	Y/N/N	Y/N/N	N/N/N	Y/Y/Y
American Society of Anesthesiologists score > 2/NNIS > 2/ /concurrent infection	Y/N/Y	N/N/N	N/N/Y	Y/N/Y	N/N/N	N/N/Y	Y/Y/Y
<b>Susceptibility to antimicrobials</b>							
meropenem/imipenem/piperacillin/tazobactam	R/R/R	S/S/S	S/R/R	R/R/R	S/S/S	R/R/R	R/R/R
amikacin/gentamicin/cefepime	S/S/R	S/S/S	R/R/R	R/R/R	S/S/S	R/R/R	R/R/R
ceftazidime/cefotaxime/ciprofloxacin	R/R/R	S/S/S	R/R/R	R/R/R	S/S/S	R/R/R	R/R/S
colistin/tigecycline	ND/ND	ND/ND	S/S	S/S	S/ND	S/S	S/ND
definitive therapy (specified)	IMP + CL	CTX + CL	CTX + CL	IMP + CL	MRP + CL	MRP + CL	IMP + CL
appropriate definitive therapy/FDTET (days)/DOEA (days)	Y/1/16	Y/5/14	Y/1/28	Y/5/15	Y/2/28	Y/3/17	Y/5/7
mortality	Y	Y	N	Y	N	Y	Y

CL – colistin, CSF – cerebrospinal fluid, CTX – ceftriaxone, DOEA – duration of effective antibiotics, F – female, FATD – from admission to diagnosis, FDTET – from diagnosis to effective treatment, IMP – imipenem-cilastatin, M – male, MRP – meropenem, N – no, ND – no available data, NNIS – National Nosocomial Infections Surveillance, R – resistant, S – susceptible, Y – yes.

in whom PCM occurred included older age (15/1,821), emergency procedure (16/2,524), CSF leakage (15/1,822), external ventricular drainage (12/1,256), intensive care unit admission (11/1,160), drain placement duration more than 72 hours (13/993), surgery duration more than 4.5 hours (6/312), repeated operations (7/565), head trauma (6/351). The pathogens isolated from the CSF were *S. aureus* in 8 (36.4%), *A. baumannii* in 7 (31.8%), *P. aeruginosa* in 2 (9.1%), *K. pneumoniae* in 2 (9.1%), *E. cloacae* in 1 (4.5%), *E. coli* in 1 (4.5%), and *E. faecalis*

in 1 (4.5%). Of the 22 patients with PCM, 14 (63.6%) had at least one concurrent HAI, and the distribution of these HAI events was SSI in 15 (68.2%), pneumonia in 4 (28.6%), bacteremia in 4 (28.6%), urinary tract infection in 2 (14.3%), and central venous catheter-related infection in 1 (7.1%). Of the 22 patients with PCM, 6 (27.3%) died. There were no significant statistical differences between patients with ABM and non-ABM (Tab. 1), except for head traumatic injury (OR = 5.873, 95% CI 1.575–17.511;  $p = 0.004$ ), and infected sites other than the central

nervous system (CNS) (OR = 0.872, 95% CI = 0.665–1.621;  $p = 0.003$ ). In this study, we disclosed that patients after traumatic head injury had higher risk of ABM (Graph 1). The clinical information of ABM is listed in Tab. 2.

## Discussion

This study showed that ABM emerged in post-traumatic head injury patients with PCM. Many studies showed that non-fermentative gram-negative bacteria (GNB), most commonly *Acinetobacter spp.*, have become important in recent years [4,8,9].

We conducted an evidence-based literature review because ABM has been rarely suggested as a cause of CNS infection [10–14]. Among the 22 PCM patients, PCM was caused by *A. baumannii* in 5 of 6 patients with traumatic head injury and in 2 of 16 patients without traumatic head injury. Among the 22 PCM patients, PCM was caused by *A. baumannii* in 6 of 8 patients with only one CNS infection source. *A. baumannii* and *S. aureus* were the most commonly isolated agents in PCM and this is in line with the suggested global emergence and increase of *A. baumannii* colonization and infection, resulting from skin or wound colonization, invasive procedures, sourced from a head traumatic wound or nostrils, and some kind of predominant cluster. In our study, *S. aureus* was still the most common pathogen in patients with PCM but we emphasize that physicians should also concentrate on *A. baumannii* when treating PCM patients.

There were no significant statistical differences between meningitis group and non-meningitis group in key risk factors. However, 5 (71.4%) of 7 ABM had a history of traumatic head injury and they died. Our results are similar to those of Baltas's report that showed 6 of 12 infectious agents being GNB among 860 patients who

developed complications after traumatic injury [9,10]. We identified eight post-craniotomy *S. aureus* meningitis and seven post-craniotomy ABMs with 6 of them suffering from traumatic head injury. Hence, we assume that patients after traumatic head injury undergoing craniotomy easily acquire *A. baumannii* and *S. aureus* infection.

In conclusion, we found a high rate of PCM due to *A. baumannii* and *S. aureus* among patients with traumatic head injury. The changing pattern of infectious agents in PCM over time suggests the need for further studies to provide the most up-to-date insight to physicians.

## References

1. Chen C, Zhang B, Yu S, et al. The incidence and risk factors of meningitis after major craniotomy in china: a retrospective cohort study. PLoS One 2014;9(7):e101961. doi: 10.1371/journal.pone.0101961.
2. McClelland S, Hall WA. Postoperative central nervous system infection: incidence and associated factors in 2111 neurosurgical procedures. Clin Infect Dis 2007;45(1):55–9.
3. van de Beek D, Drake JM, Tunkel AR. Nosocomial bacterial meningitis. N Engl J Med 2010;362(2):146–54. doi: 10.1056/NEJMra0804573.
4. Yadegarynia D, Gachkar L, Fatemi A, et al. Changing pattern of infectious agents in postneurosurgical meningitis. Caspian J Intern Med 2014;5(3):170–5.
5. Metan G, Alp E, Aygen B, et al. Carbapenem-resistant acinetobacter baumannii: an emerging threat for patients with post-neurosurgical meningitis. Int J Antimicrob Agents 2007;29(1):112–3.
6. Kourbeti IS, Vakis AF, Ziakas P, et al. Infections in patients undergoing craniotomy: risk factors associated with post-craniotomy meningitis. J Neurosurg 2015;122(5):1113–9. doi: 10.3171/2014.8.JNS132557.
7. National Nosocomial Infections Surveillance System. National Nosocomial Infections Surveillance (NNIS) System Report, data summary from January 1992 through June 2004, issued October 2004. Am J Infect Control 2004; 32(8):470–85.
8. Howard A, O'Donoghue M, Feeney A, et al. Acinetobacter baumannii: an emerging opportunistic pathogen. Virulence 2012;3(3):243–50. doi: 10.4161/viru.19700.
9. Baltas I, Tsoulfa S, Sakellariou P, et al. Posttraumatic meningitis: bacteriology, hydrocephalus, and outcome. Neurosurgery 1994;35(3):422–6.
10. Karaiskos I, Galani L, Baziaka F, et al. Successful treatment of extensively drug-resistant Acinetobacter baumannii ventriculitis and meningitis with intraventricular colistin after application of a loading dose: a case series. Int J Antimicrob Agents 2013;41:480–3. doi: 10.1016/j.ijantimicag.2013.02.010.
11. Chen CH, Chen YM, Cheng CY. Two cases reports of Acinetobacter baumannii meningitis: an emerging threat for post-neurosurgical patients: case report. J Intern Med Taiwan 2014;25:436–45.
12. Fernandez-Viladrich P, Corbella X, Corral L, et al. Successful treatment of ventriculitis due to carbapenem-resistant Acinetobacter baumannii with intraventricular colistin sulfomethate sodium. Clin Infect Dis 1999;28(4):916–7.
13. Khawcharoenporn T, Apisarnthanarak A, Mundy LM. Intrathecal colistin for drug-resistant Acinetobacter baumannii central nervous system infection: a case series and systematic review. Clin Microbiol Infect 2010;16(7): 888–94. doi: 10.1111/j.1469-0691.2009.03019.x.
14. Cascio A, Conti A, Sinardi L, et al. Post-neurosurgical multidrug-resistant Acinetobacter baumannii meningitis successfully treated with intrathecal colistin. A new case and a systematic review of the literature. Int J Infect Dis 2010;14(7):e572–9. doi: 10.1016/j.ijid.2009.06.032.

Na webových stránkách **www.csnn.eu**  
naleznete aktualizované pokyny pro autory.