

Intestinal pseudo-obstruction syndrome in a patient with acute polyradiculoneuritis

Syndrom střevní pseudoobstrukce u nemocného s akutní polyradikuloneuritidou

Dear Editor,

Acute colonic pseudo-obstruction (ACPO) is characterized by acute colonic dilatation in the absence of a mechanical obstruction. It occurs in a variety of settings, including inflammatory, neoplastic, post-traumatic, and systemic conditions. Progressive colonic distension may lead to ischaemia and subsequent perforation in approximately 15% of patients, with an associated mortality rate of up to 40% [1]. In 1948, Sir William Ogilvie described two patients with pseudo-obstruction syndrome associated with malignant infiltration of the prevertebral ganglia [2].

Autonomic innervation of the intestinal wall and its imbalance play a crucial role in the pathophysiology of intestinal pseudo-obstruction. Ogilvie originally attributed this condition to deprivation of sympathetic innervation [2]. However, current evidence suggests a predominance of sympathetic tone with reduced parasympathetic activity. Impaired parasympathetic innervation, particularly in the distal colon, results in an atonic segment and functional obstruction [3].

Among the wide range of conditions associated with intestinal pseudo-obstruction, neurological disorders such as Parkinson's disease, other neurodegenerative diseases, and drug-induced states (e.g., due to sedatives) are commonly reported [4]. Polyneuropathies are rarely implicated, and only a limited number of reports describe an association with Guillain-Barré syndrome [5]. We present a case of a 78-year-old patient with acute inflammatory demyelinating polyradiculoneuritis (AIDP) complicated by severe intestinal pseudo-obstruction.

A 78-year-old man was admitted to the Department of Neurology on November 18, 2024 with acute onset of upper and lower limbs weakness and paraesthesias of the hands and feet, which began earlier that

morning. During the day, both motor weakness and sensory disturbances progressed, predominantly in the distal extremities (tactile hypesthesia). At approximately 17:30, while attempting to walk to the toilet with forearm crutches, his legs gave way and he was unable to stand.

Neurological examination revealed moderate quadriparesis, more pronounced distally. The patient was able to hold his head elevated for up to 20 s. Grip strength measured 26 kPa in the right hand and 22 kPa in the left. In the outstretched arm position, the right arm dropped within 4 s, whereas the left was maintained for more than 20 s. Lower limb elevation was sustained for 15 s on the right and 20 s on the left. Cranial nerves were intact. Sensory examination showed impairment (tactile hypesthesia) in the distal halves of the hands and fingers and approximately 10 cm below the patella bilaterally; vibration sense was absent in the lower limbs.

Electromyography performed on November 21 confirmed a demyelinating polyradiculoneuropathy. Cerebrospinal fluid analysis demonstrated protein-cytological dissociation (protein 0.64 g/L, leukocyte count 1/μL), confirming the diagnosis of AIDP. Intravenous immunoglobulin therapy (Kiovig, total dose 2 g/kg over 5 days) was administered, resulting in mild clinical improvement. Brain CT (November 18 2024) was unremarkable; brain MRI (November 20) revealed multiple small non-expansive lesions and an arachnoid cyst in the left temporal region.

The patient was a retired clerk. His medical history included atrial fibrillation (since 2009) treated with repeated cardioversions (CHA₂DS₂-VASc score 2) and long-term anticoagulation with dabigatran. Following amiodarone therapy, he had maintained sinus rhythm since 2022. Additional comorbidities included arterial hypertension, im-

The Editorial Board declares that the manuscript met the ICMJE "uniform requirements" for biomedical papers.

Redakční rada potvrzuje, že rukopis práce splnil ICMJE kritéria pro publikace zasílané do biomedicínských časopisů.

**K. Šimková¹, L. Ungermann²,
E. Ehler^{1,3}, I. Štětkařová⁴**

¹ Department of Neurology Pardubice Regional Hospital, Czech Republic

² Radiodiagnostic Department of the Pardubice Regional Hospital, Czech Republic

³ Department of Neurology, Faculty of Health Studies, University of Pardubice, Czech Republic

⁴ Department of Neurology, Third Faculty of Medicine, Královské Vinohrady University Hospital, Prague, Czech Republic



doc. MUDr. Edvard Ehler, CSc.
Department of Neurology
Pardubice Regional Hospital
Kyjevská 44
532 03 Pardubice
e-mail: eda.ehler@tiscali.cz

Accepted for review: 13. 8. 2025

Accepted for print: 3. 6. 2026

paired glucose tolerance, dilation of the ascending aorta, and renal cysts. He had a history of COVID-19 infection in December 2021.

The clinical course became complicated on November 26, when the patient developed abdominal pain, hypogastric tenderness, and mild enterorrhagia, accompanied by mild anaemia. Abdominal ultrasound revealed meteorism and a right renal cyst. On November 29, his condition deteriorated, with subfebrile temperature, lethargy, leuco-




Fig. 1. Postcontrast CT scan with a reconstruction of 3mm layers in the coronal plane. (A) Significant dilatation of the caecum with a fluid content. (B) Dilatation of the caecum and colon sigmoideum.

C – caecum; D – colon descendens; R – rectum; S – colon sigmoideum

Obr. 1. CT vyšetření po podání kontrastní látky s rekonstrukcí v 3mm vrstvách v koronární rovině. (A) Výrazná dilatace slepého střeva s obsahem tekutiny. (B) Dilatace slepého střeva a sigmoidu.

C – slepé střevo; D – colon descendens; R – rektum; S – colon sigmoideum

cytosis (18,6), and potassium levels ranging from 3.5–4.4 mmol/L. Chest X-ray showed no signs of pneumonia but revealed elevation of the right hemidiaphragm. Oral intubation and mechanical ventilation installed.

On December 2, the patient developed diffuse abdominal pain, recurrent vomiting, abdominal distension with tympanic percussion, and reduced bowel sounds. Urgent abdominal CT demonstrated massive dilatation of the caecum and ascending colon (115 × 125 mm), elevation of the right hemidiaphragm, and compressive atelectasis of the middle and lower lobes (Fig. 1). Based on clinical and radiological findings, a diagnosis of ACPO (Ogilvie syndrome) was established. Conservative management was initiated, including neostigmine (0.5 mg s.c. every 6 h), rectal decompression, dietary restriction, and consideration of endoscopic decompression;

however, endoscopic intervention was not indicated.

By December 18, the patient was conscious, cooperative, and communicative. Abdominal findings had significantly improved. Antibiotic therapy was discontinued on December 26. Mechanical ventilation was terminated on January 1, 2025, and intensive rehabilitation was initiated. On January 30, the patient was transferred to a rehabilitation facility. At that time, he was off prokinetic therapy, tolerating a puréed diet, able to sit independently, and ambulate with a walker. Gastrointestinal function was stable.

Neurological causes account for approximately 27% of ACPO cases and include stroke, Parkinson's disease, medications (e.g., antidepressants, antipsychotics, antiparkinsonian drugs), and electrolyte disturbances such as hypokalaemia [6]. Colonic motility depends heavily on autonomic innervation,

with peristaltic activity typically originating in the caecum and propagating distally (Tab. 1). The transition zone is located near the splenic flexure, reflecting the interface between vagal and sacral parasympathetic innervation. Dysfunction of this autonomic regulation, involving the enteric nervous system and interstitial cells of Cajal, results in impaired motility [2].

Both reduced and uncoordinated autonomic stimulation may lead to severe motility disturbances. Although Guillain–Barré syndrome associated with intestinal pseudo-obstruction has been reported, it remains rare [5]. Our findings are consistent with previously published cases [7–10].

ACPO must be distinguished from paralytic ileus and mechanical obstruction. While ileus may result from vascular compromise or mechanical causes, pseudo-obstruction presents with marked colonic dilatation without an identifiable obstruction,

Tab. 1. Causes of acute colonic pseudo-obstruction.

Cardiac diseases	myocardial infarction, congestive heart failure
Surgical interventions	intra-abdominal, urological, gynaecological, orthopaedic, cardiac, neurosurgical - spine
Electrolyte disorders	hyponatremia, hypokalemia, hypocalcemia, hypercalcemia, hypomagnesemia
Metabolic disorders	diabetes mellitus, hypothyroidism, hepatopathy
Respiratory disorders	chronic obstructive pulmonary disease, mechanical ventilation
Renal disorders	renal insufficiency, nephrolithiasis
Infections	herpes zoster, sepsis, cytomegalovirus
Inflammatory causes	appendicitis, cholecystitis, pancreatitis, gastritis, pneumonia
Medication	opioids, anticholinergics, calcium channel blockers, laxatives, antidepressants, antipsychotics, epidural anaesthesia, benzodiazepines
Oncological causes	small cell lung cancer, multiple myeloma, acute myeloid leukemia, pelvic radiation, retroperitoneal invasion
Neurological causes	Parkinson's disease, dementia, stroke, multiple sclerosis, Alzheimer's disease
Old age	

potentially progressing to ischaemia and perforation.

The incidence of ACPO has decreased in recent years. Wells et al. (2017) reported approximately 100 cases per 100,000 hospital admissions, with mortality reduced to 6.4% from previously reported rates of up to 30% [2]. However, complications such as perforation or ischaemia still occur in 10–20% of cases and are associated with high mortality.

We report a rare case of AIDP complicated by severe ACPO. Early recognition and multidisciplinary management were essential for a favourable outcome. The patient survived a life-threatening condition and continues rehabilitation with gradual functional improvement.

Conflict of interest

The authors declare they have no potential conflicts of interest concerning drugs, products, or services used in the study.

Literatura

1. Underhill J, Munding E, Hayden D. Acute colonic pseudo-obstruction and volvulus: pathophysiology, evaluation, and treatment. *Clin Colon Rectal Surg* 2021; 34(4): 242–250. doi: 10.1055/s-0041-1727195.
2. Wells CI, O'Grady G, Bissett IP. Acute colonic pseudo-obstruction: a systematic review of aetiology and mechanisms. *World J Gastroenterol* 2017; 23(30): 5634–5644. doi: 10.3748/wjg.v23.i30.5634.
3. Arthur T, Burgess A. Acute colonic pseudo-obstruction. *Clin Colon Rectal Surg* 2022; 35(3): 221–226. doi: 10.1055/s-0041-1740044.
4. Vanek P, Urban O, Falt P. Percutaneous endoscopic cecostomy for management of Ogilvie's syndrome: a case series and literature Review with an update on current guideline. [online]. Available from: <https://doi.org/10.1007/s00464-023-10281-w>.

5. Pérez-Lara JL, Santana Y, Hernández-Torres J et al. Acute colonic pseudo-obstruction caused by dexmedetomidine: a case report and literature review. *Am J Case Rep* 2019; 20: 278–284. doi: 10.12659/AJCR.913645.

6. Haj M, Haj M, Rockey DC. Ogilvie's syndrome: management and outcomes. *Medicine (Baltimore)* 2018; 97(27): e11187. doi: 10.1097/MD.00000000000011187.

7. Man BL, Fu YP. Intestinal pseudo-obstruction as a presenting symptom of Guillain-Barré syndrome. *BMJ Case Rep* 2014; 2014: bcr2014205155. doi: 10.1136/bcr-2014-205155.

8. Ramirez R, Zuckerman MJ, Hejazi RA et al. Treatment of acute colonic pseudo-obstruction with tegaserod. *Am J Med Sci* 2010; 339(6): 575–576. doi: 10.1097/MAJ.0b013e3181db6b95.

9. Sahu TA. An atypical descending variant of guillain-barré syndrome with bulbar palsy, autonomic instability, and delayed colonic pseudo-obstruction: a case report. *Cureus* 2025; 17(10): e95802. doi: 10.7759/cureus.95802.

10. Luo L, Chen L, Li J et al. Case report: successful treatment of severe Guillain-Barré syndrome with paralytic ileus as a presenting symptom by intensive immunotherapy. *Front Immunol* 2025; 16: 1435817. doi: 10.3389/fimmu.2025.1435817.